



Research Article

Evaluation of the Antifungal Effects of Indole Carboxylic Acid-Pyridine Derivatives on Clinical Isolates of *Aspergillus*

Mohammad karimi¹, Mojtaba Taghizadeh Armaki², Jalal Jafarzade², Asieh Khalilpour³

1. Student Research Committee, Babol University of Medical Sciences, Babol, I.R.Iran
2. Infectious Diseases and Tropical Medicine Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R.Iran
3. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R.Iran

Article Info.

Received: 26 May 2025
Revised: 13 July 2025
Accepted: 27 July 2025
Published: 25 August 2025

* Corresponding Author:

Asieh Khalilpour
E-mail:
asieh.khalilpour@gmail.com
Tel: 011-32190101

Cite this article

karimi M, Taghizadeh Armaki M, Jafarzade J, Khalilpour A. Evaluation of the Antifungal Effects of Indole Carboxylic Acid-Pyridine Derivatives on Clinical Isolates of *Aspergillus*. Curr Res Med Sci. 2025; 8: 61-69.

Abstract


Background: Due to the increasing resistance of *Aspergillus* species to azole drugs, the development of new antifungal agents with fewer side effects and potent activity is critical. This study evaluates the antifungal activity of indole carboxylic acid-pyridine derivatives against clinical isolates of *Aspergillus*.

Methods: This cross-sectional laboratory study examined the antifungal sensitivity of 100 clinical isolates of *Aspergillus* species using the broth microdilution method (CLSI M38-A2 protocol). The minimum inhibitory concentrations (MICs) of five derivatives (K1-K5) and itraconazole were determined visually after 48 hours of incubation at 35°C. Statistical analyses were performed using SPSS version 22, and significance was set at $p < 0.05$.

Results: There was no statistically significant correlation between the MIC values of the tested derivatives and itraconazole (Wilcoxon signed-rank test, $p < 0.001$). None of the derivatives exhibited superior antifungal effects. Itraconazole consistently demonstrated lower MICs across all *Aspergillus* species compared to the tested derivatives.

Conclusion: The indole carboxylic acid-pyridine derivatives exhibited significantly weaker antifungal activity compared to itraconazole. Further structural optimization is needed to improve their efficacy.

Keywords: Indole derivatives, pyridine scaffold, itraconazole, *Aspergillus*, antifungal resistance.



Introduction

Heterocyclic compounds are among the most significant structural classes in medicinal chemistry. The indole nucleus, with the chemical formula C_8H_7N (1), is a heterocyclic scaffold fundamental to many biologically active molecules (2). To date, over ten thousand indole derivatives with biological activity have been identified, with more than 200 currently in clinical use or trials (3). They encompass various therapeutic categories, including antihistamines (4), antifungals (5), antimicrobials (6), antioxidants (7), plant growth regulators (8), anti-HIV agents (9), anticonvulsants (10), anti-inflammatory (11), and analgesics (12).

Pyridine, with the formula C_5H_5N (13), is another important heterocycle found in numerous natural products, vitamins, alkaloids, and coenzymes, as well as in many drugs and pesticides (14, 15). Recent studies suggest compounds containing the pyridine nucleus exhibit notable antibacterial, antifungal, and antiviral properties, especially when combined with other heterocycles.

Increased incidence and mortality from fungal infections, particularly among immunocompromised patients (e.g., with cancer, tuberculosis, or HIV/AIDS), underscore the need for effective antifungal agents (16-19). *Aspergillus* infections, especially invasive aspergillosis, can have mortality rates as high as 80% in some populations. Resistance to azole

drugs complicates treatment, reducing efficacy and increasing failure rates. Therefore, continuous research into heterocyclic derivatives, including indole and pyridine compounds, is essential to develop new therapies and combat drug resistance (20-23).

Despite limited local studies, the potential antifungal activity of these compounds warrants investigation. This study evaluates the antifungal effects of indole carboxylic acid-pyridine derivatives on clinical *Aspergillus* isolates collected from Babol, with the goal of informing future therapeutic strategies.

Methods

Isolates

In this experimental and laboratory study, 50 *Aspergillus* isolates: *Aspergillus fumigatus* (33), *A. flavus* (34), *A. Niger* (18), *A. terreus* (12), *A. nidulans* (3) were available in the reservebank of the Department of Parasitology and Mycology, Babol University of Medical Sciences. Fungal isolates were confirmed using the PCR–restriction fragment length polymorphism molecular technique (24). To perform the antifungal susceptibility test, fungal isolates were cultured on Sabouraud dextrose agar culture medium containing chloramphenicol and incubated for 2 to 5 days at 30 °C.

[Downloaded from crms.mubabol.ac.ir on 2026-02-19]

antifungal susceptibility tests were replicated to ensure reproducibility.

Statistical analysis

The data were analyzed using SPSS software, version 27 (IBM) software. The independent t-test was used to analyze quantitative results, and the chi-squared test was used to analyze qualitative variables, with $P \leq 0.05$ considered significant. Also, using Excel version 2020, MIC50, MIC90, and GM (geometric mean) were calculated for all isolates.

Results

Based on descriptive statistics of the frequency and percentage of the examined species among

100 *Aspergillus* isolates, the following were identified: *Aspergillus fumigatus* (33 isolates), *Aspergillus flavus* (34 isolates), *Aspergillus Niger* (18 isolates), *Aspergillus terreus* (12 isolates), and *Aspergillus nidulans* (3 isolates). In a separate analysis of each derivative compared to the drug itraconazole, the Wilcoxon test yielded a P-value of <0.001 , indicating no significant relationship between them. This suggests that the derivatives exhibit lower antifungal activity against itraconazole. Moreover, no statistically significant differences were observed in the average MIC values of the derivatives when compared to one another (see Tables 1 and 2).

Table 1: Results of the Wilcoxon Signed-Rank Test

Sample	MIC ($\mu\text{g/ml}$)	MIC Control ($\mu\text{g/ml}$)	P-value
K1	356.90 $\pm$ 476.80	0.179042 $\pm$ 0.22235	<0.001
K2	361.06 $\pm$ 480.00	0.179042 $\pm$ 0.22235	<0.001
K3	286.85 $\pm$ 417.92	0.179042 $\pm$ 0.22235	<0.001
K4	308.44 $\pm$ 423.68	0.179042 $\pm$ 0.22235	<0.001
K5	311.53 $\pm$ 440.96	0.179042 $\pm$ 0.22235	<0.001

Table 2: Descriptive statistics of drug sensitivity results

Species	K1	K2	K3	K4	K5	Itraconazole
<i>Aspergillus fumigatus</i>						
Minimum	64	128	64	128	128	0.031
Maximum	1024	1024	1024	1024	1024	0.5

MIC	256	512	512	256	512	0.125
(µg/ml)50						
MIC (µg/ml)	1024	1024	1024	1024	1024	0.5
90						
GM	309.27	441.99	381.55	381.25	381.55	0.132
<i>Aspergillus</i>						
<i>flavus</i>						
Minimum	128	64	64	64	64	0.031
Maximum	1024	1024	1024	1024	1024	0.5
MIC 50	512	256	256	512	256	0.25
MIC	1024	1024	1024	1024	1024	0.5
(µg/ml)50						
MIC (µg/ml)	400.88	326/.95	307.307	326.95	298/309	0.155
90						
<i>Aspergillus</i>						
<i>niger</i>						
Minimum	128	128	256	64	64	0.031
Maximum	1024	1024	1024	1024	1024	0.5
MIC	512	256	256	512	256	0.25
(µg/ml)50						
MIC (µg/ml)	1024	1024	1024	1024	1024	0.5
90						
GM	335/20	348/36	376/25	310/35	322/53	0.16
<i>Aspergillus</i>						
<i>terreus</i>						
Minimum	128	64	64	128	256	0.031
Maximum	1024	1024	512	1024	1024	0.5
MIC	256	256	256	256	512	0.125
(µg/ml)50						
MIC (µg/ml)	512	512	512	1024	512	0.25
90						
GM	304.43	241.63	203.18	287/35	406/37	0.11
<i>Aspergillus</i>						
<i>nidulans1</i>						
<i>Aspergillus</i>	256	512	128	256	128	0.125
<i>nidulans2</i>						
<i>Aspergillus</i>	1024	256	1024	512	512	0.25
<i>nidulans3</i>						

- **MIC 50:** Minimum Inhibitory Concentration at which 50% of the organisms are inhibited
- **MIC 90:** Minimum Inhibitory Concentration at which 90% of the organisms are inhibited
- **GM:** Geometric Mean

Discussion

Fungal pathogens are responsible for a wide spectrum of diseases, ranging from life-threatening invasive infections such as fungemia, meningitis, and pneumonia, to chronic conditions including allergic bronchopulmonary aspergillosis and chronic pulmonary aspergillosis. A significant proportion of these infections arise in immunocompromised individuals, where the associated mortality rates are alarmingly high. Although early diagnosis and administration of antifungal therapy are critical to clinical outcomes, the current arsenal of antifungal agent's remains limited (26). Moreover, the rise of antifungal resistance, particularly among *Aspergillus* species, has created a pressing need for the development of new antifungal compounds with novel mechanisms of action (27).

Heterocyclic compounds, especially those containing indole and pyridine rings, have drawn significant attention in drug development due to their broad-spectrum biological activities. These scaffolds are commonly found in therapeutic agents with antifungal, antibacterial, antiviral, anti-inflammatory, and anticancer properties.

Previous studies have demonstrated the antifungal potential of indole-based compounds. For instance, Vaca et al. (2020) evaluated several indole alkaloid derivatives and reported that compounds bearing free amino groups showed potent antifungal activity against

phytopathogenic fungi, such as *Moniliophthora Roreri* (28). Similarly, Shirinzadeh et al. (2018) synthesized indole derivatives substituted with triazole, thiadiazole, and carbothioamide moieties, and found that compound 3d (an indole-triazole) exhibited superior activity against *Candida krusei* compared to fluconazole, largely due to the presence of electron-withdrawing substituents like m-chlorophenyl (29).

Kokorekin and colleagues developed aryl thiocyanates derived from indole frameworks and demonstrated their strong antifungal activity against *Candida albicans*, *Candida krusei*, and *Aspergillus Niger* (MIC = 0.12 µg/mL). Their findings indicated that small changes in side chains, such as replacing a methyl nitrogen group with a phenyl ring, could drastically affect antifungal potency. These results collectively emphasize that subtle structural variations—particularly in the substitution pattern of the indole nucleus—play a critical role in determining antifungal efficacy (30).

In the current study, we synthesized a series of indole carboxylic acid–pyridine derivatives through a multi-step process. This involved the initial formation of 4-(4-aminophenylamino)-N-methylpicolinamide via the reaction of 2-picolinic acid with thionyl chloride and methylamine, followed by condensation with 4-amino-N-methylbenzamide. Subsequent alkylation using benzyl chloride derivatives and coupling with 2-chloroacetamide analogs yielded

the final target compounds. The structural integrity of intermediates was confirmed, and all reaction stages were monitored using thin-layer chromatography.

Despite successful synthesis, the antifungal evaluation revealed that the tested derivatives (K1–K5) exhibited significantly higher MIC values ($\geq 256 \mu\text{g/mL}$) than itraconazole ($\text{MIC}_{50} = 0.125 \mu\text{g/mL}$), indicating markedly lower antifungal potency. Statistical analysis confirmed this finding ($p < 0.001$). These results are in contrast to previous reports on indole-based compounds with improved efficacy. The observed discrepancy may be attributed to differences in substitution groups (e.g., absence of thiocyanate or halogenated moieties), the nature of the linker units, and the overall hydrophobicity or steric hindrance imposed by the side chains.

Furthermore, variations in the synthetic pathway, such as solvent-free heating conditions and non-functionalized terminal moieties, might have influenced the bioactivity of the final compounds. It is also possible that the electronic and spatial orientation of the carboxylic acid–pyridine fragment relative to the indole core did not favor optimal interaction with fungal cell targets.

Taken together, while the current derivatives were structurally inspired by known bioactive scaffolds, their limited antifungal activity suggests the need for further structure–activity

relationship (SAR) optimization. Future studies should focus on introducing electron-withdrawing substituents (e.g., halogens, nitro groups), exploring triazole or thiocyanate hybrids, and modifying the central scaffold to enhance membrane permeability and binding affinity.

Conclusion

This cross-sectional study examined the prevalence of internet addiction among high school students in Babol, Iran, and revealed that 27.9% exhibited moderate levels of internet addiction. The results indicate significant associations between internet addiction and factors such as gender, educational field, and family structure, with female students and those in technical fields showing higher rates of moderate addiction. The study also underscores the importance of the home environment and parental employment status in influencing students' internet usage and potential addiction. Understanding these characteristics is crucial for developing targeted interventions aimed at reducing internet addiction among adolescents by addressing specific demographic influences.

Acknowledgment

This research was conducted with financial support and is the result of a research project with an ethical code (IR.MUBABOL.REC.1401.068). We would like to express our sincere gratitude to all those who helped us in this research.

Conflicts of interest: There are no conflicts of interest.

Availability of data and material: Not applicable.

Author's contribution

Study concept and design: RH, SJ, MEH, and HS. Analysis and interpretation of data: RH, MEH, SH. Critical revision of the manuscript RH, MEH, SSM. Data collection: MAL, Data collection: SSM, Writing an article: MEH. RH. All the authors had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Consent for publication: Not applicable.

References

1. Khalilpour A, Asghari S. Synthesis, characterization and evaluation of cytotoxic and antioxidant activities of dihydropyrimidone substituted pyrrole derivatives. *Medicinal Chemistry Research*. 2018 Jan;27:15-22.
2. Ghashari K, Mahdavi Omran S, Khalilpur A, Jafarzadeh J, Taghizadeh Armaki M, Hossein Nejad A, Aminian AR. Antifungal Susceptibility of 3, 4-di-hydropyrimidine-1-(H2)-L-H1-pyrrole Derivatives in Candida Clinical Isolates. *Journal of Mazandaran University of Medical Sciences*. 2022 Dec 10;32(215):26-34.
3. Bronner SM, Im GYJ, Garg NK. *Indoles and indolizidines*. Wiley-VCH: Weinheim, Germany; 2011. p. 221-65.
4. Battaglia S, Boldrini E, Da Settimo F, Dondio G, La Motta C, Marini AM, et al. Indole amide derivatives: synthesis, structure-activity relationships and molecular modelling studies of a new series of histamine H1-receptor antagonists. *European journal of medicinal chemistry*. 1999;34(2):93-105.
5. Przheval'Skii N, Magedov I, Drozd V. New derivatives of indole. Synthesis of s-(indolyl-3) diethyl dithiocarbamates. *Chemistry of Heterocyclic Compounds*. 1997;33(12):1475-6.
6. Al-Hiari YM, Qaisi AM, El-Abadelah MM, Voelter W. Synthesis and antibacterial activity of some substituted 3-(aryl)-and 3-(heteroaryl) indoles. *Monatshefte für Chemie/Chemical Monthly*. 2006;137(2):243-8.
7. Tan D-X. Melatonin: a potent, endogenous hydroxyl radical scavenger. *Endocr j*. 1993;1:57-60.
8. Abele E, Abele R, Dzenitis O, Lukevics E. Indole and Isatin Oximes: Synthesis, Reactions, and Biological Activity .*Chemistry of Heterocyclic Compounds*. 2003;39(1):3-35.
9. Suzen S, Buyukbingol E. Evaluation of anti-HIV activity of 5-(2-phenyl-3'-indolal)-2-thiohydantoin. *Il Farmaco*. 1998;53(7):525-7.
10. El-Gendy AA, Said MM, Ghareb N, Mostafa YM, El-Ashry ESH. Synthesis and Biological Activity of Functionalized Indole-2-carboxylates, Triazino-and Pyridazino-indoles. *Archiv der Pharmazie: An International Journal Pharmaceutical and Medicinal Chemistry*. 2008;341(5):294-300.
11. Kumar A, Sharma S, Malik N, Sharma P, Kaushik K, Saxena KK, et al. Synthesis of anti-inflammatory, analgesic and COX-II inhibitory activities of indolylpyrazolines. 2004.
12. Smith AL, Stevenson GI, Swain CJ, Castro J. Traceless solid phase synthesis of 2, 3-disubstituted indoles. *Tetrahedron letters*. 1998;39(45):8317-20.
13. Khalilpour A, Asghari S, Pourshab M. Synthesis and characterization of novel Thiazolo [3, 2-a] pyrimidine derivatives and evaluation of antioxidant and cytotoxic activities. *Chemistry & Biodiversity*. 2019 May;16(5):e1800563.
14. Hamada Y. Role of pyridines in medicinal chemistry and design of BACE1 inhibitors possessing a pyridine scaffold. *Pyridine*. 2018 Jul 18:9-26.
15. Ling Y, Hao Z-Y, Liang D, Zhang C-L, Liu Y-F, Wang Y. The expanding role of pyridine and dihydropyridine scaffolds in drug design. *Drug Design, Development and Therapy*. 2021;15:4289.
16. Altaf AA, Shahzad A, Gul Z, Rasool N, Badshah A, Lal B, et al. A review on the medicinal importance of pyridine derivatives. *J Drug Des Med Chem*. 2015;1(1):1-11.
17. Zalaru C, Dumitrascu F, Draghici C, Tarcomnicu I, Tatia R, Moldovan L, et al. Synthesis, spectroscopic characterization, DFT study and antimicrobial activity of novel alkylaminopyrazole derivatives. *Journal of Molecular Structure*. 2018;1156:12-21.

18. Marinescu M, Cintează LO, Marton GI, Chifiriuc M-C, Popa M, Stănculescu I, et al. Synthesis, density functional theory study and in vitro antimicrobial evaluation of new benzimidazole Mannich bases. *BMC chemistry*. 2020;14(1):1-16.
19. Marinescu M. Synthesis of antimicrobial benzimidazole-pyrazole compounds and their biological activities. *Antibiotics*. 2021;10(8):1002.
20. Zilberberg MD, Harrington R, Spalding JR, Shorr AF. Burden of hospitalizations over time with invasive aspergillosis in the United States, 2004–2013. *BMC Public Health*. 2019;19(1):1-7.
21. Ostrosky-Zeichner L, Al-Obaidi M. Invasive fungal infections in the intensive care unit. *Infectious Disease Clinics*. 2017;31(3):475-87.
22. Taccone F, Van den Abeele A, Bulpa P, Misset B, Meersseman W, Cardoso T. Paiva 346 JA, Blasco-Navalpotro M, De Laere E, Dimopoulos G, Rello J, Vogelaers D, Blot SI, Asp 347 ICUSI (2015) Epidemiology of invasive aspergillosis in critically ill patients: clinical 348 presentation, underlying conditions, and outcomes. *Critical care (London, England)*. 19(1):7.
23. Bassetti M, Carnelutti A, Righi E. Issues in the management of invasive pulmonary aspergillosis in non-neutropenic patients in the intensive care unit: A role for isavuconazole. *IDCases*. 2018;12:7-9.
24. Ardebilifard A, Hoseinnejad A, Jafarzade J, Hajizadeh Jouybari N, Rajabnia Baboli A, Khalilpour A. Investigating the Antifungal Effects of Spirocyclopropane Oxindoles Derivatives Against *Aspergillus* Species. *Research in Molecular Medicine*. 2023; 11 (2): 103-112.
25. Heydari A, Taghizadeh Armaki M, Jafarzade J, Vahidi B, Yazdizadeh M, Babaii N, Khalilpour A. Evaluation of the Efficacy of Synthetic Indole-Hydrazono-Thiazolidinone Compounds on *Candida* Isolates from Oral Infections. *Journal of Mazandaran University of Medical Sciences*. 2024 Oct 10;34(237):30-9.
26. Parslow BY, Thornton CR. Continuing shifts in epidemiology and antifungal susceptibility highlight the need for improved disease management of invasive candidiasis. *Microorganisms*. 2022 Jun 13;10(6):1208.
27. Fisher MC, Alastruey-Izquierdo A, Berman J, Bicanic T, Bignell EM, Bowyer P, Bromley M, Brüggemann R, Garber G, Cornely OA, Gurr SJ. Tackling the emerging threat of antifungal resistance to human health. *Nature reviews microbiology*. 2022 Sep;20(9):557-71.
28. Vaca J, Salazar F, Ortiz A, Sansinenea E. Indole alkaloid derivatives as building blocks of natural products from *Bacillus thuringiensis* and *Bacillus velezensis* and their antibacterial and antifungal activity study. *The Journal of Antibiotics*. 2020;73(11):798-802.
29. Shirinzadeh H, Süzen S, Altanlar N, Westwell AD. Antimicrobial activities of new indole derivatives containing 1, 2, 4-triazole, 1, 3, 4-thiadiazole and carbothioamide. *Turkish Journal of Pharmaceutical Sciences*. 2018;15(3):291.
30. Kokorekin V, Terent'Ev A, Ramenskaya G, Grammatikova N, Rodionova G, Ilovaiskii A. Synthesis and antifungal activity of arylthiocyanates. *Pharmaceutical Chemistry Journal*. 2013;47(8):422-5.