

# Current Research in Medical Sciences

ISSN: 2538-6212



# Stress, anxiety, depression among nurses caring for COVID-19 patients in Babol, Iran: a logistic regression



# Parvin Aziznejadroshan <sup>1</sup>, Fatemeh Shaker Zavardehi <sup>2</sup>, Mojtaba Qanbari Qalehsari <sup>3</sup>

- 1. Non-Communicable Pediatric Disease Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran.
- 2. B.S.C. Laboratory Science, Department of Pathology, Shahid Beheshti Hospital, Babol University of Medical Sciences, Babol, Iran.
- 3. Department of Nursing, Faculty of Nursing and Midwifery, Babol University of Medical Sciences, Babol, Iran.

#### **Article Info**

#### ABSTRACT

# Article type: Research Article

**Background:** Nurses who are involved in the caring of COVID-19 patients, are at risk of mental distress. The present study was conducted with the purpose of demonstrating the relationship between stress, anxiety, depression and characteristics of nurses who provide care to COVID-19 patients.

**Methods:** This descriptive cross-sectional study was conducted using the census method from May to June, 2020 on 224 clinical nurses who were working in hospitals affiliated with Babol University of Medical Sciences and were involved in caring for COVID-19 patients. The data collection instruments included the Depression, Anxiety and Stress Scale - 21 items (DASS-21) and the individual characteristic questionnaire. Bivariate and multivariate binary logistic regressions were computed to identify the associated factors.

Results: The mean and the standard deviation for stress, anxiety and depression scores were 9.47±7.30, 9.29±7.51 and 8.84±7.22 respectively. 17.4% had stress, 54% had anxiety and 43% had a degree of depression. There were significant relationships between the nurses' stress level and characteristics including age (OR =3.009, 95%CI  $1.46-6.16,\ P=0.003),\ having\ children\ (OR=0.26,\ 95\%CI\ 0.11-0.63,\ P=0.003),\ work$ experience (OR=4.50, 95% CI 2.17-9.96, P=0.000) and employment status (OR=0.39, 95% CI 0.16-0.95, P=0.04). Moreover, along with these characteristics, job satisfaction (OR =3.03, 95%CI 5.64-1.63, P=0.000), level of physical activities (OR =0.26, 95% CI 0.08–0.82, P=0.02), exercising (OR =2.27, 95% CI 1.31-3.90, P=0.003) and violence in the workplace (OR =0.27, 95%CI 0.12- 0.56, P=0.001) also had significant relationships with the nurses' anxiety level. Furthermore, the relationships between the nurses' level of depression and characteristics including age (OR =2.07, 95%CI 1.15 - 3.72, P=0.014), work experience (OR =21.79, 95%CI 1.04 - 3.10, P=0.03), job satisfaction (OR=3.03, 95%CI 1.63-5.64, P=0.000), exercising (OR =1.76, 95% CI 1.02-3.04, P=0.04), having chronic diseases (OR =0.35, 95% CI 0.15-0.81, P=0.014), violence in the workplace (OR =0.39, 95% CI 0.20–0.75, P=0.005) and sleep (OR =1.77, 95%CI 1.00–3.16, P=0.050) were significant.

**Conclusion:** The authorities should consider a number of the individual characteristics of nurses including age, work experience, gender, marital status, having children, job satisfaction, sleep, violence in the workplace and history of chronic disease in their planning and provide psychological support for them. Psychological support services may be needed for nurses in order to protect and promote their mental health.

Received: 12 June 2023

**Accepted:** 11 September 2023

**Published:** 16 September 2023 **Keywords:** Stress, Anxiety, Depression, COVID-19, Nursing care, Nurses.

**Cite this article:** Aziznejadroshan P, et al. Stress, anxiety, depression among nurses caring for COVID-19 patients in babol, Iran: a logistic regression. *Current Research in Medical Sciences*. 2023; 7(2): 1-12.

© The Author(s).

Publisher: Babol University of Medical Sciences

Address: Department of Nursing, Faculty of Nursing and Midwifery, Babol University of Medical Sciences, Babol, Iran.

**Tel:** +98 11 32190597 **E-mail:** m2ghanbari@yahoo.com

<sup>\*</sup> Corresponding Author: Mojtaba Qanbari Qalehsari

# Introduction

The COVID-19 pandemic is a serious threat for public health (1). Nurses work at the frontline of the fight against this disease and risk their lives while performing their duties (2). They have always played a key role in preventing and controlling infection, care and public health (3). Early in the spread of COVID-19, more than 3000 medical staff in Wuhan, China, got infected (40% in hospitals and 60% in the society) (3).

In the face of this critical situation, the healthcare workers at the forefront, who are directly involved in the diagnosis, treatment and caring of COVID-19 patients, are at risk of mental distress and other mental health disorders. The increasing number of confirmed and suspected cases, the high level of work pressure, the lack of personal protective equipment, the extensive media coverage, the shortage of specific drugs and feeling that they are not being sufficiently supported may all affect the healthcare workers' stress (4). A whole new working condition, the job burn-out caused by the burden of heavy workload and the protective equipment, fear of getting infected and contaminating others, a feeling of inability to attend to clients and Maintaining proper communication in stressful situations are the challenges faced by the health care providers (5).

The emergency situations caused by COVID-19 put major pressures on nursing services. When nurses work in environments where job demands are high and resources are low, they are exposed to higher job stress and more physical and psychological symptoms, which might have negative effects on their health and well-being (6, 7). Maintaining the mental health of the nursing staff is necessary to control infectious diseases (8).

Evidence has shown that while nurses are caring for patients with emerging diseases such as MERS, Covid-19 and H1N1, they experience loneliness, anxiety, fear, fatigue, sleep disorders and other mental and physical problems (9,10,11). A review study showed that demographic variables including gender, occupation, age, workplace and department as well as psychological variables such as self-efficacy and poor social support were accompanied by increase in stress, anxiety, depression symptoms and insomnia in healthcare workers. Evidences indicated that COVID-19 can be an independent risk factor for stress in healthcare workers (12). In a research in 2020 in China on mental health of healthcare workers who were exposed to COVID-19, the findings showed that 50.4% of them reported depression symptoms, 44.5% anxiety, 34% sleep disorder and 71.5% symptoms of distress (13). The prevalence rates of moderate to severe depression, anxiety, and stress among Iranian nurses who provided care to patients with COVID-19 were 43.7%, 73%, and 24%, respectively (14).

The findings of another study showed that the mean self-reported stress and anxiety scores were  $32.19\pm7.56$  and  $39.91\pm12.92$  respectively and the multiple regression analysis showed that having a child in the family, working hours per week and anxiety were the main variables related to the nurses' stress (3).

Since COVID-19 is an emerging disease and the healthcare system and culture vary from region to region, more studies is needed on the psychological experience of frontline nurses coping with COVID-19 (15). Regular screening of healthcare workers involved in patient care as well as diagnosing and providing care for patients with COVID-19 to assess their stress, depression and anxiety seems necessary (12). The existing information about work stress among nurses is limited (3).

Considering the important role of nurses in improving and promoting healthcare services in the society, the researchers decided to investigate the relationship between stress, anxiety, depression and some of the individual characteristics of nurses involved in caring for COVID-19 patients in hospitals affiliated with Babol University of Medical Sciences.

#### **Methods**

This descriptive cross-sectional study was conducted from May to July, 2020 using census method on all clinical nurses working in two hospitals affiliated with BUMS (Ayatollah Rouhani and Shahid Yahya Nejad) in northern Iran who were involved in direct care of COVID-19 patients. Among the 302 nurses who directly involved in the care of Covid-19 patients, 224 completed the questionnaire. The inclusion criteria were: having a bachelor's or master's degree in nursing, being directly involved in providing care for COVID-19 patients in one of the wards of the two aforementioned hospitals, at least one-year experience of working in the hospital and willingness to participate in the study. Participants who incompletely answered the study instruments were excluded. This study was approved by the university research ethics committee with the code of ethics of IR.MUBABOL.HRI.REC.1399.138.

#### **Data Collection Tools**

The research instruments used in this study included the individual characteristics questionnaire and the short form version of Depression Anxiety Stress Scales (DASS-21). The individual characteristics questionnaire had 18 items comprising age, gender, marital status, education level, work experience, type of work shift, job satisfaction, average amount of overtime, employment status, types of hospital, having children, level of physical activity, exercising, history of chronic diseases, workplace violence, sleep and history of depression in the family. DASS-21 is a self-report instrument that shows the recent mood swings and was first introduced by Lovibond in 1995(13). As the name suggests, it has 21 items which are scored on Likert scales. These 21 items contain: 7 questions related to stress including finding it hard to wind down, tendency to over-react to situations, using a lot of nervous energy, getting agitated, finding it difficult to relax, being intolerant of anything that keeps you from getting on with what you were doing, and feeling touchy; 7 questions related to anxiety including dryness of the mouth, breathing difficulty, trembling, being worried about situations in which you might panic and make a fool of yourself, feeling close to panic, action of the heart in the absence of physical exertion, and feeling scared without any good reason; and finally 7 questions related to depression including inability to experience any positive feeling at all, finding it difficult to work up the initiative to do things, feeling that you have nothing to look forward to, feeling down-hearted and blue, being unable to become enthusiastic about anything, feeling that you are not worth much as a person, and feeling that life is meaningless. The validity and reliability of this instrument have been reviewed and approved by Mehdipour and Najafi (16, 17). The internal consistency of the scale was determined in the study of Shamsaie et al. (2018) using Cronbach's alpha and the figures for each subscale were as follows: 85% for stress, 86% for anxiety and 83% for depression (18).

#### Data collection

After defining the purpose of the study and reassuring the participants that all information will remain confidential, the researcher distributed the questionnaires and gave explanation in case any of the items were unclear. The participants had to read each statement and rate the intensity of the symptom mentioned in it on a 4-point Likert scale as it had been experienced over the past week. In this regard, zero meant that the symptom had not been experienced at all, one meant it had been experienced to some extent, two meant it had been experienced rather much, and three meant it had been experienced very much. The total score of each subscale of this instrument is obtained by summing up the scores of the questions related to that subscale. Since DASS-21 is the short-form version of the main scale (42-item), the total score of each subscale must be doubled (18). Then, the intensity of the signs and symptoms can be determined according to table1.

Symptom severity of unferent parameters in the time				
Score variables	stress	anxiety	depression	
normal	0-14	0-7	0-9	
mild	15-18	8-9	10-13	
moderate	19-25	10-14	14-20	
severe	26-33	15-19	21-27	
Extremely severe	>34	>20	>28	

Table 1. Symptom severity of different parameters in the three groups

Lovibond, S.H. & Lovibond, P.F. (1995). Manual for the Depression Anxiety & Stress Scales. (2nd Ed.) Sydney: Psychology Foundation.

#### Data Analysis

Data were analyzed using SPSS statistical software version 18. The correlations between independent and dependent variables were tested by binary logistic regression analysis. The variables which had a p-value less than 0.2 were entered into multivariate analysis. In the end, the model was finalized using backward elimination. Finally, a p-value of less than 0.05 was considered statistically significant association, and the adjusted odds ratio with 95% CI was calculated to determine the strength of association.

Regular monitoring of data collection was performed by the principal investigator. The completed questioners were cheeked manually first for completeness then the collected data were entered onto the computer. If a questionnaire is incomplete, it will be returned for completion, the questionnaire was excluded from the analysis if they did not wish to fill in the missing information. The executive stages of the study were approved by the Vice Chancellor for Research of Babol University of Medical Sciences. Informed consent for voluntary participation in this study was obtained from all participants.

#### **Results**

The study sample contained 224 nurses with an average age of 32.9±6.31. The demographic characteristics of the participants were as follows: 167 (74.6%) were female, 148 (68%) were younger than 50 years of age, 85% had bachelor's degree, 80% worked rotating shifts, 58% had 1 to 10 years of work experience, 67% were working under a formal contract, 80% were married, 59% had at least one child, 52% were working 50 to 100 hours of overtime per month, 67% said they were satisfied with their jobs, 90% had physical activities, 53% did not exercise regularly, 87% did not have any chronic diseases, 98% did not have any history of mental disorders, 90% had no history of depression in their families, 68% got 7 to 8 hours of sleep at least 3 to 4 days a week, and 78% did not experience workplace violence (Table 2).

# Level of Stress, Anxiety and Depression

The mean ( $\pm$  standard deviation) stress score calculated by DASS-21 was 4.73 ( $\pm$ 3.65), which was multiplied by two (to make it equal to the result of the main scale) and the result was 9.47 ( $\pm$ 7.30). More than 82% of the nurses had normal stress levels. The mean ( $\pm$  standard deviation) anxiety score was 4.67 ( $\pm$ 3.75) based on DASS-21 and 9.34 ( $\pm$ 7.51) after multiplying it by two. More than half of the nurses (54%) had degrees of anxiety. The mean ( $\pm$  standard deviation) depression score was 4.42 ( $\pm$ 3.61) as measured by DASS-21 and 8.84 ( $\pm$ 7.22) after it was doubled. About 43% of the nurses experienced some degrees of depression (Table 3).

Table 2. Frequency distribution of respondents by socio-demographic characteristics and selected variables

selected variables					
Demographic-social variables		Number (%)			
Sex	male	57(25.4)			
SEA	female	167(76.4)			
999	≤35	148(68)			
age	>35	68(32)			
	morning	23(10.3)			
Shift type	Evening or night	22(9.8)			
	rotational	178(79.8)			
Words Ermanian on	≤10	129(57.6)			
Work Experience	>10	95(42.5)			
	bachelor of Nursing	191(85.3)			
<b>Education level</b>	Master of Nursing	33(14.7)			
Differential 3	single	43(19.2)			
Marital status	married	181(80.8)			
W-2 191	Yes	132(59.5)			
Having a child	no	90(40.5)			
T. 1	Yes	149(67)			
<b>Employment status</b>	no	71(33)			
	50<	38(17)			
0 (1 (1)	50-100	116(51.8)			
Overtime(h)	100-150	60(26.8			
	>150	10(4.5)			
	Rouhani Hospital	141(62.9)			
hospital	Yahya Nejad Hospital	83(37.1)			
Tab madisferation	Yes	149(67.7)			
Job satisfaction	no	71(32.3)			
DI 1 1 11 11 1	inactive	20(8.9)			
Physical activity level	active	203(91.1)			
	yes	105(47.1)			
exercise	no	118(52.9)			
C1	yes	28(12.50			
Chronic disease	no	195(87.5)			
¥¥73 1 * 1	yes	48(21.6)			
Workplace violence	no	174(78.4)			
Matria 7 Ob also 2 Att	yes	151(68)			
Maintain 7–8 h sleep 3–4 times a week	no	71(32)			
D	yes	5(2.2)			
Psychiatric disorder	no	218(97.8			
Formally, 12.44 6.3	yes	22(9.9)			
Family history of depression	no	200(90.1)			

Table 3. Frequency	distribution of nu	recel depression	anvioty and atreas
Table 3. Frequency	distribution of nu	rses' denression.	anxiety and stress

z usto et z requesto,	y distribution of nurses depression, anxiety and stre			
Score variables	stress	anxiety	depression	
	Number (%)	Number (%)	Number (%)	
normal	181 (82)	100 (45.7)	127 (57.7)	
mild	15 (6.7)	26 (11.9)	42 (19.1)	
moderate	17 (7.6)	44 (20.1)	36 (16.4)	
severe	4 (1.8)	27 (12.3)	12 (5.58)	
Very severe	3 (1.3)	22 (10)	3 (1.4)	

# Multivariate Analyses

In the final model and in the presence of all variables, five variables including marital status, work experience, having children, job satisfaction and history of depression emerged as the significant stress-related factors. Among these factors, work experience was the strongest variable related to stress in nurses involved with COVID-19, in a way that the nurses who had more than 10 years of experience had a 4 times greater risk of experiencing stress than those nurses who had less than 10 years of experience (OR=4.5).

Moreover, in the final model and in the presence of all variables, three variables including work experience, job satisfaction and violence in the workplace emerged as the factors associated with anxiety in nurses involved with COVID-19. Among these factors, again work experience had the strongest relationship with anxiety in nurses, in a way that the nurses who had more than 10 years of experience were 3.87 times as likely to experience anxiety as the nurses who had less than 10 years of experience (aOR= 3.87, 95% CI=2.09-7.18).

Finally, regarding the factors associated with depression, six variables including gender, history of depression, exercising, sleep pattern, history of chronic diseases and violence in the workplace emerged in the final model and in the presence of all variables. The nurses who did not have a history of depression had 0.27 times greater risk of depression than the nurses with a history of depression and the nurses who did not exercise were 2.04 times as likely to suffer from depression as the nurses who did exercise. Moreover, the nurses who got 7 to 8 hours of sleep less than 3 to 4 days a week were 2.28 times as likely to get depressed as the nurses who had a normal sleep pattern and the nurses who did not experience violence in their workplaces and had no history of chronic diseases had respectively 0.39 and 0.32 times greater risk of depression than the nurses who experienced workplace violence and had chronic diseases (Table 4).

Table 4: Multiple logistic regression model predicting stress, anxiety and depression in nurses involved with Covid-19

variables	groups	SE	P-Value	AOR	95% Conf. Interval
stress					
constant		20.27	0.37	8.42	0.07-942.42
Marital status	single	-	-	-	-
	married	0.15	0.03	0.21	0.04-0.90
Years of employment	<10	-	-	-	-
	≥11	1.98	0.001	4.50	1.89-10.69

variables	groups	SE	P-Value	AOR	95% Conf. Interval		
Having a child	yes	-	-	-	-		
	no	0.11	0.009	0.16	0.04-0.63		
Job satisfaction	yes	ı	=	ı	-		
	no	1.08	0.01	2.66	1.19-5.90		
Depression history	yes	-	-	-	-		
Depression instory	no	0.17	0.003	0.31	0.10-0.93		
	Anxiety						
constant		0.42	0.39	0.42	0.05-3.07		
work experience	<10	-	-	-	-		
Work experience	≥11	1.22	0.000	3.87	2.09-7.18		
Job satisfaction	yes	-	-	-	-		
<b>0</b> 00 5 <b>00</b> 2	no	0.92	0.004	2.67	1.36-5.25		
Workplace violence	yes	-	-	-	-		
Troringiace Holenee	no	0.12	0.003	0.30	0.13-0.67		
	Depre	ession					
constant	-	193.08	0.005	114.02	4.10-3151.12		
sex	male	-	-	-	-		
	female	0.20	0.10	0.55	0.26-1.14		
Depression history	yes	-	-	-	-		
Dopt of motory	no	0.13	0.01	0.27	0.10-0.73		
exercise	yes	-	-	-	-		
	no	0.65	0.02	2.04	1.08-3.83		
Maintain 7–8 h sleep 3–4 times a week	yes	-	-	-	-		
- Tancou week	no	0.73	0.01	2.28	1.22-4.27		
Chronic disease	yes	-	-	-	-		
	no	0.14	0.01	0.32	0.13-0.79		
Workplace violence	yes	-	-	-	-		
workplace violence	no	0.13	0.009	0.39	0.19-0.78		

# **Discussion**

The results showed that 17.4% of the nurses had stress, more than half (54%) experienced anxiety and about 43% had some degrees of depression. In the study of Lai et al. (2020) on the healthcare workers who were involved in providing care for COVID-19 patients in China, 50.4% of them had symptoms of depression, 44.5% anxiety, 34% sleep disorder, and 71.5% distress (4). The study of Gupta et al. (2020) in Nepal indicated that 38% of the healthcare workers involved in caring for COVID-19 patients suffered from anxiety or depression (19). A meta-analysis showed that the prevalence of anxiety and depression in healthcare workers who provide care for COVID-19 patients was 23.2% and 22.8% respectively (20). Other studies showed that the prevalence of anxiety is about 11.3 to 50% (4, 21, 22). The prevalence rates of moderate to severe depression, anxiety, and stress among Iranian nurses who provided care to patients with COVID-19 were 43.7%, 73%, and 24%, respectively (14). This is in line with the findings of several previous studies which reported that the prevalence of stress was 48% among nurses in Iran during the COVID-19 pandemic (23). The findings of the present study report that it is slightly higher. The high prevalence of anxiety in Iranian nurses could be related to the lack of protective equipment and fear of infection. Previous studies have also confirmed the high levels of anxiety in those who had direct clinical contact with COVID-19 patients (19, 22).

In this study, prevalence of depression was 43%, which is higher than what was reported in other studies in other region of the world. In this regard, we can refer to a study in Nepal that reported a prevalence of depression of 8% (18) and 28% in a study in China (22). Moreover, in a meta-analysis, the combined prevalence was reported to be 22.8% (20). The high prevalence of depression in Iranian nurses is probably related to risk factors such as the high transmission risk of COVID-19 and the social isolation of nurses due to concern about spreading the disease to relatives and friends.

In this study, 17.4% of the nurses experienced some degrees of stress, which is similar to the results of the study of Wu et al. (24) However, in the study of Mo et al. (2020) the mean stress score of Chinese nurses was reported to be 39.9%. The long work hours during a week as well as fear of getting infected by COVID-19 infection through the respiratory droplets and direct contact (3) increased the Chinese nurses' stress to a relatively higher level compared to Iranian nurses. Generally, stress in health care workers fighting COVID-19 is attributed to long shifts, unrealistic wages, lack of personal protective equipment, and concern about getting infected or infecting one's family (25, 26).

Work experience was the strongest variable associated with stress and anxiety among nurses who were involved with COVID-19 in a way that nurses who had more than 10 years of experience were at an increased risk of experiencing stress and anxiety compared to the nurses with less than 10 years of experience. Long work hours during a week is one of the factors affecting nurses' stress and anxiety (3). The Iranian nurses who were satisfied with their jobs reported less stress, anxiety and depression than those who were not interested in and satisfied with their jobs. In the study of Letavak et al. (2012) it was perceived that the nurses who have a lower level of job satisfaction get more depressed (27). There are other studies that show job satisfaction and professional commitment help to reduce stress and anxiety (28, 29). Moreover, a history of depression emerged as a factor associated with stress and depression. Many of the previous studies also showed that high stress levels in nurses lead to anxiety, frustration, depression and other mental and emotional disorders (30, 31).

Marital status and having a child in the family were among factors affecting the nurses' stress. Having children causes stress in nurses because the children need care and also because there is a risk of transmission of COVID-19 from the nurse parent to the child, which is in line with the study of Mo (3). Moreover, there was a significant relationship between marital status and nurses' stress, which is consistent with the study of Yildirim et al. (2020) (32).

The nurses who are older than 35 years are more likely to get stress and anxiety. Although in the study of Yildirim et al. (2020), it was shown that the young healthcare staff had higher anxiety scores compared to older ones (32). In addition, the nurses who are informally employed are less likely to

experience stress and anxiety than the nurses who are formally employed. The type of employment is associated with workplace anxiety (33). It seems that informally employed nurses can quit their jobs more easily because they have no obligation to the organization. As a result, they have less stress and anxiety than formally employed nurses.

Violence in the workplace emerged as one of the factors associated with stress and anxiety in nurses involved with COVID-19. In the study of Tong et al. (2019), it was found that workplace violence was followed by symptoms of depression in nurses. Reducing the workplace violence and developing the psychological capital can be helpful in the fight against the symptoms of depression (34).

The nurses who exercise and have higher levels of physical activities are less likely to get depression than the nurses who do not exercise and have low levels of physical activities. Physical inactivity and sedentary behavior are significantly associated symptoms of depression and anxiety (35).

The risk of depression is higher in nurses who have a chronic disease or an abnormal sleep pattern. The study of Levatak et al. (2012) indicates that the nurses who have health problems are more likely to be depressed. In other words, there is a relationship between nurses' medical problems and their depression (27). The study of Wu showed that more than one third of the medical staff had symptoms of insomnia during the COVID-19 outbreak, which was associated to isolation and the psychological distress caused by the prevalence of the disease. Considering various psychological factors, it is necessary for the medical personnel to receive interventions in this regard (36). The distress caused by symptoms of anxiety affects the medical staff's sleep quality (37).

The nurses who fight against COVID-19 are generally under psychological pressure. The authorities should consider the nurses' individual characteristics (age, work experience, gender, marital status, having children and history of chronic diseases), improve their working conditions and provide opportunity for them to get enough sleep to help them modulate their stress, anxiety and depression. Moreover, the nurses should be taught appropriate sports and physical activities to do in their leisure time as well as relaxation techniques to maintain their mental health.

# **Conclusion**

Prevalence of stress, anxiety and depression is high among nurses involved with caring for COVID-19 patients and it is associated with a number of their individual characteristics including age, work experience, gender, marital status, having children, job satisfaction, workplace violence, sleep, exercising and history of chronic diseases. Therefore, screening services and psychological supports (comprehensive interventions for the early diagnosis and management of the symptoms of depression, anxiety, and stress also education about stress management) should be provided for nurses and their working conditions should be improved to reduce their anxiety and distress.

# Limitations

The present study has several limitations. This study examined only the nurses in the city of Babol in North of Iran. Therefore, it is difficult to generalize the results to all Iranian nurses. Since this was a cross-sectional study, it was not possible to observe the subjects in a longer period of time like a longitudinal study. Third, only one questionnaire was used in this study because of the time constraints.

**Acknowledgement:** The authors would like to thank the Research Deputy of Babol University of Medical Sciences, nursing managers and nurses of Ayatollah Rouhani and Shahid Yahya Nejad Hospitals who helped us in this research.

**Authors' Contributions:** Conceptualization: P.A. and F.S.Z; Methodology: P.A., F.S.Z and M.Q.Q; Statistical analysis and investigation: M. Q.Q.; Writing - original draft preparation: P.A. and F.S.Z.; Writing - review and editing: P.A. and M.Q.Q; Supervision: P.A.

**Funding:** This research received a grant from Babol University of Medical Sciences (Code: 9909619). The Babol University of Medical Sciences did not have any role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript

**Competing Interests:** The authors declare no conflicts of interest.

**Data availability:** The data that supports the findings of this study are available from the corresponding author upon reasonable request.

### References

- 1. Pan, Y., Guan, H., Zhou, S., Wang, Y., Li, Q., Zhu, T. Xia, L. Initial CT findings and temporal changes in patients with the novel coronavirus pneumonia (2019-nCoV): A study of 63 patients in Wuhan, China European Radiology.2020; https://doi.org/10.1007s 00330-020-06731-x.
- 2. Catton, H., RN, MA, BS (Econ) (Hons). (2020). Global challenges in health and health care for nurses and midwives everywhere. International Nursing Review, 2020; 67(1): 4–6.
- 3. Mo Y, Deng L, Zhang L, Lang Q, Liao C, Wang N, Qin M, Huang H. Work stress among Chinese nurses to support Wuhan in fighting against COVID-19 epidemic. J Nurs Manag. 2020; 28(5): 1002–1009. https://doi.org/10.1111/jonm.13014
- 4. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, Wu J & etal .Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. JAMA Network Open. 2020; 3(3): e203976. https://doi.org/10.1001/jamanetworkopen.2020.3976.
- 5. Liu Q, Luo D, Haase JE, Guo Q, Wang X Q, Liu S, Xia L, Liu Z & etal. The experiences of health-care providers during the COVID-19 crisis in China: a qualitative study. www. thelancet. com/lancetgh Vol 8 June 2020.
- 6. Chou, L. P., Li, C. Y., & Hu, S. C. Job stress and burnout in hospital employees: Comparisons of different medical professions, in a regional hospital in Taiwan. British Medical Journal Open.2014: 4 ,e004185.
- 7. Khamisa, N., Oldenburg, B., Peltzer, K., Ilic, D. Work related stress, burnout, job satisfaction and general health of nurses International Journal of Environmental Research and Public Health.2015; 12(1): 652–666. https://doi.org/10.3390/ijerp h1201 00652.
- 8. Kang, L., Li, Y. I., Shaohua, H. U., Chen, M., Yang, C., Yang, B. X., ...Liu ,Z. The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus. Lancet Psychiatry. 2020; 7: e14.
- 9. Kim Y. Nurses' experiences of care for patients with Middle East respiratory syndrome coronavirus in South Korea. Am J Infect Control. 2018; 46:781–787.
- 10. Khalid I, Khalid TJ, Qabajah MR, Barnard AG, Qushmaq IA. Healthcare worker semotions, perceived stressors and coping strategies during MERS-CoV outbreak. Clin Med Res. 2016; 1303:1–22.
- 11. Honey M, Wang WY. New Zealand nurse's perceptions of caring for patients with influenza A (H1N1). Nurs Crit Care. 2013;18: 63–69.
- 12. Spoorthya MS, Pratapa S K, Mahant S. Mental health problems faced by healthcare workers due to the COVID-19 pandemic–A review. Asian Journal of Psychiatry. 51 (2020) 102119. https://doi.org/10.1016/j.ajp.2020.102119
- 13. Lovibond PF, Lovibond SH. The structure of negative emotional states comparison of the depression anxiety stress scales (Dass) with the beck depression and anxiety inventories. Behav Res Ther. 1995; 33:335–43.

- 14. Sharifi A, Fallahi-Khoshknab M, Mohammadi S, Zeraati M, Jamshidi Z, Aghabeygi-Arani M, Mirzaei N, Fallahi-Khoshknab N, Rasooli P. Depression, anxiety, and stress among Iranian nurses in COVID-19 care wards. BMC Psychol. 2022; 20;10(1):205. doi: 10.1186/s40359-022-00911-8.
- 15. Sun N, Wei L, Shi S, Jiao D, Song R, & etal .A qualitative study on the psychological experience of caregivers of COVID-19 patients. American Journal of Infection Control. 2020; 48: 592–598.
- 16. Mehdipour-Rabori R, Nematollahi M. The effect of recommended Azkar on anxiety, stress, and depression in families of patients undergoing open heart surgery. Iran J Nurs Midwifery Res. 2014;19(3):238-41. pmid: 24949060
- 17. Najafi Kalyani M, Pourjam E, Jamshidi N, Karimi S, Kalyani VN. Survey of stress, anxiety, depression and self-concept of students of Fasa University of medical sciences, 2010. J Fasa Univ Med Sci. 2013; 3(3):235-40.
- Shamsaei F, Yaghmaei S, Sadeghian E, Tapak L. Survey of Stress, Anxiety and Depression in Undergraduate Nursing Students of Hamadan University of Medical Sciences. Iranian Journal of Psychiatric Nursing (IJPN). 2018; 6(3):26-30. DOI: 10.21859/ijpn-06034
- 19. Gupta A K, Mehra A, Niraula A, Kafle K, Deo S P, Singh B, Sahoo S, Grover S. Prevalence of anxiety and depression among the healthcare workers in Nepal during the COVID-19 pandemic. Asian Journal of Psychiatry. 2020; 54: 102260. doi: 10.1016/j.ajp.2020.102260
- 20. Pappa S, Ntella V, Giannakas T, Giannakoulis V.G, Papoutsi E, Katsaounou P. Prevalence of depression, anxiety, and insomnia among health care workers during the COVID-19 pandemic: a systematic review and meta-analysis. Brain Behav Immun. 2020.https://doi.org/10.1016/j.bbi. 2020.05.026. S0889-1591(20)30845-X. Advance online publication
- 21. Zhu J., Sun L., Zhang L., Wang H., FanA., Yang B., Li W., Xiao S. Prevalence and influencing factors of anxiety and depression symptoms in the first-line medical staff fighting against COVID-19 in Gansu. Front. Psychiatry.202; 11, 386. https://doi.org/10.3389/fpsyt.2020.00386
- 22. Liu, C.Y., Yang, Y.Z., Zhang, X.M., Xu, X., Dou, Q.L., Zhang, W.W., Cheng, A., 2020. The prevalence and influencing factors in anxiety in medical workers fighting COVID-19 in China: a cross-sectional survey. Epidemiol. Infect. 148, e98. https://doi.org/10. 1017/S0950268820001107.
- 23. Sarboozi Hosein Abadi T, Askari M, Miri K, Nia MN. Depression, stress and anxiety of nurses in COVID-19 pandemic in Nohe-Dey Hospital in Torbat-e-Heydariyeh city, Iran. J Mil Med. 2020;22(6):526–33.
- 24. Wu, J., Wu, X., Wu, F., Dia, Y., Dechun, C. & Gong, X. Survey of sleep quality of clinic-al front-line nurses and its influencing factors, in the fight against new coronavirus pneumonia. Nursing Research. 2020; 344, 558-562.
- 25. Grover, S., Sahoo, S., Mehra, A., Avasthi, A., Subramanyan, A., 2020. Psychological impact of COVID-19 lockdown: an online survey from India. Indian J. Psychiatry
- 26. Zhang, W.R., Wang, K., Yin, L., Zhao, W.F., Xue, Q., Peng, M., Min, B.Q., Tian, Q., Leng, H.X., Du, J.L., Chang, H., Yang, Y., Li, W., Shangguan, F.F., Yan, T.Y., Dong, H.Q, Han, Y., Wang, Y.P., Cosci, F., Wang, H.X., 2020. Mental health and psychosocial problems of medical health workers during the COVID-19 epidemic in China. Psychother. Psychosom. 1–9. https://doi.org/10.1159/000507639. Advance online publication.
- 27. Letvak S, Ruhm CJ, Mccoy T. Depression in hospital- employed nurses. Clin Nurse Spec. 2012; 26(3):177-82. doi: 10.1097/NUR.0b013e3182503ef0.
- 28. Hegney DG, Craigie M, Hemsworth D, Osseiran Moisson R, Aoun S, Francis K, et al. Compassion satisfaction, compassion fatigue, anxiety, depression and stress in registered nurses in Australia: study 1 results. J Nurs Manag. 2014; 22 (4):506-18.
- 29. Khodadadi E, Hosseinzadeh M, Roghaye Azimzadeh R, Fooladi M. The relation of depression, anxiety and stress with personal characteristics of nurses in hospitals of Tabriz, Iran. International Journal of Medical Research & Health Sciences. 2016; 5(5):140-148.

- 30. Malinauskiene, V., Leisyte, P., Romualdas, M., & Kirtiklyte, K. Associations between self-rated health and psychosocial conditions Life style factors and health resources among hospital nurses in Lithuania. Journal of Advanced Nursing. 2011; 67(11), 2383–2393.https://doi.org/10.1111/j.1365-2648.2011.05685.x
- 31. Teles, M. A. B., Barbosa, M. R., Vargas, A. M. D., Gomes, V. E., Ferreira, E. F., Martins, A. M. E.B., ... Ferreira, R. C. Psychosocial work conditions and quality of life among primary health care employees. A cross sectional study. Health Quality of Life Outcomes. 2014: 12, 72. https://doi.org/10.1186/1477-7525-12-72
- 32. Yildirim TT, Atas O, Asafov A, Yildirim K, Balibey H. J Coll. Physicians Surg Pak. 2020; 30(6):26-31.
- 33. Choi E S, Jung H S, Kim S H, Park H .The influence of workplace violence on work-related anxiety and depression experience among Korean employees. J Korean Acad Nurs. 2010; 40(5):650-61. doi: 10.4040/jkan.2010.40.5.650.
- 34. Tong C, Cui C, Li Y, Wang L. The effect of workplace violence on depressive symptoms and the mediating role of psychological capital in chinese township general practitioners and nurses: A cross-sectional study. Psychiatry Investig. 2019 Dec; 16(12): 896–903.
- 35. Bélair M A, Kohen D E, Kingsbury M, Colman I. Relationship between leisure time physical activity, sedentary behavior and symptoms of depression and anxiety: evidence from a population based sample of Canadian adolescents. BMJ Open. 2018; 8: e021119. doi:10.1136/bmjopen-2017-021119
- 36. Wu J, Wu X, Wu F, Dia Y, Dechun C,ong X. Survey of sleep quality of clinic-al front-line nurses and its influencing factors, in the fight against new coronavirus pneumonia. Nursing Research. 2020; 344, 558–562. Retrieved from http://kns.cnki.net/kcms/detai l/14.1272. r.20200 214.1136.004.html
- 37. Kirwan M, Pickett SM, Jarrett NL. Emotion regulation as a moderator between anxiety symptoms and insomnia symptom severity. Psychiatry Res. 2017; 254: 40–7. doi: 10.1016/j.psychres.2017.04.028.